

KP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
12 April 2001 (12.04.2001)

PCT

(10) International Publication Number  
**WO 01/24818 A1**

- (51) International Patent Classification<sup>1</sup>: A61K 39/00, 35/14, C12N 5/08
- (74) Agent: PRINS, A., W.; Vereenigde, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).
- (21) International Application Number: PCT/NL00/00712
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 4 October 2000 (04.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/157,442 4 October 1999 (04.10.1999) US
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (for all designated States except US): LEIDS UNIVERSITAIR MEDISCH CENTRUM [NL/NL]; Albusdreef 2, NL-2333 ZA Leiden (NL).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): REA, Delphine, Gabrielle, Josette [NL/NL]; Floresstraat 1, NL-2315 HP Leiden (NL). MELIEF, Cornelis, Johannes, Maria [NL/NL]; Wilhelminapark 33, NL-2012 KC Haarlem (NL). OFFRINGA, Rienk [NL/NL]; Stieltjesstraat 63, NL-2313 SJ Leiden (NL).

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/24818 A1

(54) Title: DENDRITIC CELLS ACTIVATED IN THE PRESENCE OF GLUCOCORTICOID HORMONES ARE CAPABLE OF SUPPRESSING ANTIGEN-SPECIFIC T CELL RESPONSES

(57) Abstract: The present invention provides novel methods for immunotherapy. The invention provides immune cells and methods to generate them, with the capacity to at least in part reduce an immune response in a host. In one aspect the invention provides a method for generating a dendritic cell with the capacity to tolerate a T cell for antigen said T cell was specific for, comprising culturing peripheral blood monocytes from an individual to differentiate into dendritic cells, activating said dendritic cells in the presence of a glucocorticoid hormone and loading said activated dendritic cell with said antigen T cell was specific for.

Title: Dendritic cell activated in the presence of glucocorticoid hormones are capable of suppressing antigen-specific T cell responses.

The invention relates to the field of medicine. More in particular the invention relates to the field of immunotherapy.

5 Background of the invention

The remarkable immunostimulatory properties of DC reside in their ability to transport antigens from peripheral tissues to lymphoid organs where they present these antigens to T  
10 cells in an optimal costimulatory context (1). To achieve this complex sequence of events, DC exist in different functional stages. Immature DC behave as sentinels in peripheral tissues where they efficiently capture antigens. Upon pathogen invasion, induction of protective T cell  
15 responses requires the activation of immature DC into mature immunostimulatory cells. DC activation is triggered in inflamed tissues by cytokines such as IL-1 and TNF- $\alpha$  and by bacterial components such as LPS (2, 3). Activated DC migrate to T cell areas in the lymph nodes while upregulating their  
20 costimulatory capacities and optimizing their antigen presenting functions. Upon interaction with antigen-specific T cells, DC activation is further completed through engagement of the receptor-ligand (L) pair CD40-CD40L, leading to the production of IL-12 (4, 5, 6), a key cytokine  
25 for T helper (Th) type 1 and cytotoxic T lymphocyte (CTL) priming (7).

APC activation through CD40-CD40L interactions represents an important immunoregulatory step for the establishment of protective T cell immunity against pathogens  
30 and tumors (8, 9, 10). This process also plays a key role in the onset of destructive T cell-mediated disorders such as auto-immune diseases, allograft rejection and graft versus host disease (11, 12, 13). The current treatment of these

disorders largely relies on the administration of glucocorticoids (GC), which exert potent anti-inflammatory and immunosuppressive effects. Because GC negatively interfere with many aspects of T cell activation such as IL-2-driven proliferation and inflammatory cytokine production (reviewed in 14), activated T cells have long been considered as the main targets for GC action. Several lines of evidence now suggest a role for DC in GC-induced immune suppression. Moser et al (15) found that GC prevented the spontaneous activation of murine DC thereby decreasing their T cell stimulatory potential. Kitajima et al (16) showed that GC could hamper the T cell-mediated activation of a murine DC line. Viera et al reported that human DC exposed to GC were poor producers of IL-12 upon LPS stimulation (17). These findings only concern loss of typical DC features and therefore favor a simple inhibitory role of GC on DC activation. A more complex immunoregulatory action on the DC system has not been considered.

The present invention resulted from a detailed analysis of the impact of GC on the CD40-mediated activation of monocyte-derived DC. These DC develop after culture with GM-CSF and IL-4 (2, 18) or after transmigration through endothelial cells (19) and are known to mature into the most potent human Th1-type-inducing APC upon CD40 ligation (5, 20). Moreover, these APC can easily be generated in large numbers and are thereby the cells of choice for DC-based modulation of T cell immunity (21, 22). In contrast to previous studies, the present invention shows that GC such as dexamethasone (DEX) do not merely prohibit DC activation, but that it converts CD40 ligation on human monocyte-derived DC is transformed into an alternative activation pathway. DEX profoundly affect the CD40-dependent maturation of human monocyte-derived DC, not only by preventing the upregulation of costimulatory, adhesion and MHC surface molecules, but also by causing these cells to secrete the anti-inflammatory mediator IL-10 instead of the Th1 stimulatory cytokine IL-12. In agreement with

these phenotypic and functional changes, DC triggered through CD40 in the presence of DEX are poor stimulators of Th1-type responses. Most importantly, the present invention shows that such DC are able to induce a state of hyporesponsiveness in Th1 cells, indicating that these cells are capable of active suppression of Th1-type immunity.

As already mentioned above, the impact of GC on DC has been the subject of several previous studies by others. However, in contrast with the present invention, these studies only highlighted inhibitory effects of GC on the DC system. DEX was found to block the upregulation of CD80, CD86 and MHC class II molecules upon activation of murine spleen DC (15, 16), whereas very recently DEX was demonstrated to also prevent the differentiation of DC from monocyte precursors (28). In these studies, the inability of DC to acquire high expression of costimulatory and MHC molecules was accompanied with a decrease in their T cell stimulatory potential, but the effect of GC on IL-12 production was not investigated. On the other hand, Viera et al found that the effect of GC on LPS-induced DC activation consisted in a 4-fold reduction of IL-12p70 synthesis (17). This partial effect on IL-12 secretion contrasts with the complete suppression of IL-12p70 production which is subject of the present invention, and can be explained by the fact that their GC-treated immature DC were extensively washed prior to LPS stimulation. We indeed found that upon removal of GC, the effects of these drugs on immature DC were rapidly reversible. The continuous presence of GC during CD40 triggering of DC was clearly preferred in order to stably and completely modulate DC activation (data not shown). Taken together, previous findings indicated that the impact of GC on the DC system should be merely interpreted as an inhibitory event. Importantly, the present invention clearly demonstrates that GC such as DEX do not simply suppress DC activation but rather redirect this process towards a distinct functional program.

DC activation through engagement of CD40-CD40L is a key stimulatory event for the generation of effective Th1 and CD4-dependent CTL responses in vivo (10, 36, 37, 38). This pathway however is also involved in the development of unwanted T cell responses leading to autoimmune disease or organ-transplant rejection (11, 12, 13). Until now, treatment of patients suffering from such disorders largely relies on the systemic administration of GC hormones. This treatment does not only suppress pathogenic T cell responses but also induces a general state of immunosuppression and metabolic and endocrine side effects. The present invention demonstrates that activation of human monocyte-derived DC through CD40, in the presence of GC such as DEX, results in an IL-10-producing APC that is a poor stimulator for Th1-type responses and that can even confer hyporesponsiveness to Th1 cells. The present invention therefore indicates that such DC loaded with appropriate antigens can be exploited as a novel approach for specifically downregulating unwanted T cell responses in vivo.

The dendritic cells of the invention possess different capabilities than previously reported for dendritic cells. One can therefore consider these cells to be part of a class of cells distinct from the class formed by the "classical" dendritic cells. The dendritic cells of the invention can be used in a different way than the classical dendritic cells. The dendritic cells of the invention can for instance be used to suppress, at least in part, an undesired immune response in a host. In one aspect invention therefore provides a method for preparing a pharmaceutical composition for reducing an unwanted T cell response in a host, comprising culturing peripheral blood monocytes from said host to differentiate into dendritic cells, activating said dendritic cells in the presence of a glucocorticoid hormone and loading

said activated dendritic cells with an antigen against which said T cell response is to be reduced. An unwanted T cell response can be any type of T cell response. For instance, but not limited to; a T cell response associated with an  
5 auto-immune disease or a transplantation disease such as a graft versus host disease or a host versus graft disease. A pharmaceutical composition of the invention typically comprises a dendritic cell of the invention suspended in a liquid suitable for preserving the function of said dendritic  
10 cell in said liquid and/or suitable for administration to a host. A host preferably is a human. Preferably said host is at risk of developing or is suffering from an auto-immune disease or allergy. Preferably, said host suffers from or is at risk of suffering from host versus graft disease and/or  
15 graft versus host disease. With the term at risk is meant that one expects that said host may develop said disease, for instance but not limited to a host receiving a transplant. Such a host is considered to at risk of developing host versus graft disease. An antigen typically is a peptide  
20 capable of binding to a major histocompatibility complex I and/or II molecule. Such peptides are known in the art and a person skilled in the art is capable of determining whether a given peptide comprises an antigen or not. An antigen may be derived from a naturally occurring protein. An antigen may  
25 also be a synthetic peptide or equivalent thereof, preferably with an amino-acid sequence equivalent to a peptide derived from a protein.

In another aspect the invention provides a pharmaceutical composition for reducing an unwanted T cell response in a  
30 host, said composition being obtained by culturing peripheral blood monocytes from said host to differentiate into dendritic cells, activating said dendritic cells in the presence of a glucocorticoid hormone and loading said activated dendritic cells with an antigen against which said  
35 T cell response is to be reduced. In one embodiment a method is provided for reducing an unwanted T cell response in a

host, comprising administering a composition of the invention to said host.

The invention further provides method for reducing an unwanted T cell response in a host comprising culturing  
5 peripheral blood monocytes from said host to differentiate into dendritic cells, activating said dendritic cells and/or their precursors in the presence of a glucocorticoid hormone and loading said activated dendritic cells with an antigen against which said T cell response is to be reduced and  
10 administering said composition to said host.

In one embodiment of the invention said activation is done through a CD40 receptor. Activation of DC through triggering of the CD40 receptor can involve either incubation with a CD8-CD40L fusion protein, a trimeric form of CD40L consisting  
15 of CD40L-molecules to which a modified leucine zipper has been attached, anti-CD40 antibodies, or cells that express CD40L. Other signals that can be employed for the activation of DC as described in the present invention include lipopolysaccharide (LPS) and polyI/C.

20 In another aspect the invention provides a method for obtaining an dendritic cell capable of tolerising a T-cell for an antigen comprising providing said dendritic cell with a glucocorticoid hormone, activating said dendritic cell and providing said dendritic cell with said antigen. With the  
25 term tolerising is meant that said dendritic cell has an immunosuppressive effect on said T cell. A tolerised T cell will essentially not respond with cell division when exposed to a cell presenting an antigen said T cell would in the untolerised state respond to with cell division. A tolerised  
30 T cell will essentially not respond with killing a cell presenting an antigen said T cell would in the untolerised state respond to with cell kill.

In one embodiment said dendritic cell and/or a precursor thereof is provided with said glucocorticoid hormone in  
35 vitro. A T cell of the invention is preferably an antigen specific T cell, preferably a cytotoxic T cell or a Th cell.

In another aspect the invention provides an isolated dendritic cell capable of modifying the function of an antigen specific Th cell, which would otherwise enhance a given immune response, resulting in a T cell that is capable of reducing this immune response. In one embodiment the invention provides a method for modifying an antigen specific T-cell comprising providing an dendritic cell according to the invention with said antigen and co-cultivating said T-cell and said dendritic cell. Preferably, said co-cultivating is performed in vitro. Said method may further comprise multiplying said functionally modified T-cell. The invention also provides an isolated functionally modified T-cell obtainable by a method according to the invention.

In another aspect the invention provides the use of a glucocorticoid hormone for obtaining an dendritic cell capable of functionally modifying a T-cell. The invention also provides a pharmaceutical composition comprising an dendritic cell and/or a functionally modified T-cell according to the invention. The invention further provides the use of a dendritic cell and/or a functionally modified T-cell according the invention for the preparation of a medicament.

The invention also provides a method for the treatment of an individual suffering from or at risk of suffering from a disease associated with at least part of the immune system of said individual comprising providing said individual with an dendritic cell and/or a functionally modified T-cell according to the invention. Preferably, said dendritic cell and/or said functionally modified T-cell, or precursors thereof are derived from an HLA-matched donor. Preferably, said HLA-matched donor is said individual.

Method of treatments of the invention are preferably use for the treatment of an individual suffering from an auto-immune disease, an allergy, a graft versus host disease and/or a host versus graft disease.



## Examples

### Example 1

#### *Impairment of CD40-CD40L-mediated phenotypic changes by DEX*

5 We explored the impact of DEX on the phenotypic changes induced by CD40 ligation on immature monocyte-derived DC. In the absence of DEX, the fusion protein CD8-CD40L induced a strong upregulation of the costimulatory molecules CD80, CD86 and CD40, of the MHC class I and II molecules, of the  
10 adhesion markers CD54 and CD58 and of the DC maturation marker CD83 (Fig 1). In the presence of DEX, these CD8-CD40L-induced phenotypic changes were dramatically impaired: the upregulation of CD80, CD86, CD40, CD54, CD58 and of the MHC class I and II molecules was largely inhibited and CD83 was  
15 not expressed (Fig 1). Importantly, DEX-treated DC did not revert to a monocyte/macrophage stage as shown by the lack of expression of CD14 (Fig 1). Titration of DEX showed a complete inhibition of CD40-mediated phenotypic changes at  $10^{-6}$  M and  $10^{-7}$  M, a partial blockade at  $10^{-8}$  M and no effect  
20 at  $10^{-9}$  M and  $10^{-10}$  (data not shown). In addition, DEX action was dependent on binding to the GC-receptor, since it was abolished by simultaneous addition of the GC receptor antagonist RU486 (data not shown). In experiments performed with LPS or TNF- $\alpha$  as activation agents, similar results were  
25 obtained. However, the combination of DEX and TNF- $\alpha$  induced a massive cell death (viable cell recovery 5-10% of control cultures), a phenomenon that was not observed when DEX-treated DC were stimulated with LPS or through CD40 (viable cell recovery 60 to 100% of control cultures) (not  
30 shown).

We next analyzed whether activated DC could still be affected by DEX. DC incubated with CD8-CD40L for 48h and further exposed to DEX maintained a stable activated phenotype (Fig 2).

We conclude that DEX prevents the phenotypic changes induced by CD40 signals on immature DC and that already activated DC are resistant to DEX action.

5 Example 2

*DEX does not interfere with the regulation of DC antigen uptake machinery*

Unlike activated DC, immature DC efficiently internalize antigens through macropinocytosis and mannose receptor-mediated endocytosis (2, 3, 25, 26). We analyzed whether DEX could affect the DC antigen capture machinery and its downregulation following CD40 cross-linking. As shown in Fig 3, incorporation of FITC-BSA and FITC-mannosylated BSA by immature DC and by DEX-treated immature DC was comparable. 10 Upon CD40 triggering, a similar decrease of FITC-BSA and FITC-mannosylated BSA uptake by both DEX-treated and untreated DC was observed (Fig 3). These results were the first to indicate to us that DEX does not block all aspects of DC activation, since it does not interfere with the down-regulation of the DC antigen capture machinery. 20

Example 3

*DEX-treated CD40-triggered DC secrete IL-10 instead of IL-12*

A key feature of CD40-triggered DC for initiating T cell immunity resides in their ability to produce the proinflammatory cytokine IL-12 (5, 6, 27). We investigated whether DEX affected IL-12 production by DC stimulated through CD40, and we explored the possibility that DEX could promote the secretion of the anti-inflammatory cytokine IL-10. As shown in Fig 4, CD40 triggering of DC strongly induced IL-12p40 and IL-12p70 secretion (up to 120ng/ml and 170pg/ml respectively) but only poorly stimulated the production of IL-10 (up to 68pg/ml). In contrast, CD40 triggering of DEX-treated DC resulted in a dramatically reduced IL-12p40 30 production (up to 100 fold) and in the complete suppression of IL-12p70 secretion, whereas IL-10 production was strongly 35

enhanced (up to 50 fold) (Fig 4). Immature DC and their DEX-treated counterparts failed to secrete detectable amounts of IL-12 and IL-10 (Fig 4). Therefore, CD40 ligation of DC in the presence of DEX triggers the secretion of high levels of the anti-inflammatory cytokine IL-10 instead of IL-12.

#### Example 4

*DEX-treated CD40-triggered DC are capable of suppressing Th1-type immunity*

The strikingly modified response of DC to CD40 ligation in the presence of DEX prompted us to compare the T cell stimulatory potential of these cells with that of their DEX-untreated counterparts. In an allogeneic MLR, CD40-triggered DC induced a strong proliferative T cell response whereas the addition of DEX prior to CD40 triggering reduced their T cell stimulatory capacity to that of immature DC (Fig 5). When tested for their ability to stimulate an hsp65-specific CD4<sup>+</sup> Th1 clone, CD40-triggered DC pulsed with the hsp65 protein or with the specific peptide epitope p3-13 were found to be potent inducers of both T cell proliferation and T-cell dependent IFN-g production (Fig 5). In contrast, in the presence of Ag-pulsed DEX-treated CD40-triggered DC, T cell proliferation and IFN-g production were significantly decreased ( $p < 0.001$  and  $p < 0.01$  respectively) (Fig 5). We next investigated whether DEX-treated CD40-triggered DC were simply poor stimulators of Th1 cells, or whether they could exert suppressive effects on these T cells. We therefore tested hsp65-specific T cells stimulated with p3-13-pulsed DEX-treated CD40-triggered DC for their capacity to respond to a second potent antigenic challenge. Fig 6 shows that preculturing T cells with CD40-triggered DC led to a strong T cell proliferation and IFN-gamma production upon second antigen-specific restimulation. In contrast, preculture with DEX-treated CD40-triggered DC resulted in a dramatically reduced proliferative and IFN-gamma production capacity of Th1 cells. Thus, CD40 triggering of DC in the presence of DEX

results in APC that are not merely poor inducers of T cell responses but that also induce a state of hyporesponsiveness in Th1 cells.

5    Example 5

DEX-treated activated DC suppress anti-transplant immunity in vivo

In a first attempt to test whether DEX-treated DC could  
10 suppress unwanted immunity, a skin transplant model was set up in mice. Skin transplants from C57Bl/6 mice, when grafted onto fully allogeneic Balb/c recipient mice, are known to be rapidly rejected as a result of a very strong allo-reactive anti-transplant T cell immune response. Experiments were  
15 performed to analyse whether pre-treatment of the recipient mice with DEX-treated DC resulted in suppression of this host versus graft response. We made use of an established murine DC cell line of C57Bl/6 origin, named D1, which exhibits a stable immature phenotype in vitro unless they receive an activating signal through agonistic anti-CD40 antibodies or  
20 LPS. This trigger results in fully matured DC that are highly capable of CTL priming in vitro as well as in vivo (39;40). Maturation of D1 involves strong upregulation of the cell surface expression of the costimulatory molecule CD86 (B7.2),  
25 the CD40-receptor and Class II MHC (fig. 7). Similar to our observations in human DC, maturation of D1 in the presence of DEX largely prevented the upregulation of these molecules at the cell surface (Fig. 7). This suggested that the DEX-treated D1 cells could be exploited to suppress T cell  
30 immunity in vivo. Balb/c mice were therefore injected with D1 cells that had been activated in the absence or presence of DEX. One week after immunisation, the allo-reactive response of splenocytes of these mice, as well as of control mice, was tested in vitro. Mice pre-treated with immature D1 cells that  
35 had been cultured in the presence or absence of DEX did not exhibit significantly altered Th1-type alloreactive immunity

as compared to control mice (Fig. 8). This is in accordance with our observations with human DC, which also showed that DEX-treated immature DC did not exhibit different properties than normal immature DC. As expected, mice immunised with fully matured D1 showed enhanced responsiveness. Importantly, mice pre-treated with D1 activated in the presence of DEX displayed greatly decreased Th1-type allo-reactive immunity (fig. 8).

It was subsequently tested whether the reduction of allo-responses in the latter group of mice, as measured in vitro, would reflect decreased allo-reactive immunity in vivo. Groups of Balb/c mice were immunized with D1 cells matured in the absence or presence of DEX. Skin transplants of C57Bl/6 origin were grafted onto these mice one week later, and graft survival was monitored. The grafts were rapidly rejected in control mice and mice immunized with normally matured D1 cells (Fig. 9). Rejection was not enhanced in the latter group, despite the stronger allo-reactive responses in these mice (Fig. 8). Most likely, the response in the control mice is already sufficient for maximal rejection. Importantly, graft rejection is significantly ( $p < 0.05$ ) delayed in mice pre-treated with D1 matured in the presence of DEX (fig. 9). This outcome corresponds with the greatly reduced allo-reactivity in these mice as determined in vitro (Fig. 8).

In conclusion, the skin grafting experiments in mice demonstrate that DC matured in the presence of DEX can indeed be employed to suppress unwanted immunity such as against transplants. Although in the experiments shown the transplants are eventually rejected, it should be noted that donor and recipient displayed large differences with respect to transplantation antigens. In humans, donors and recipients are matched as closely as possible for these antigens. Murine experiments exploiting better matched donor and recipient strains are underway. The data in Fig. 8 and 9 show that pre-treatment with DEX-treated mature DC will result in much more profound effects in a this setting.

### Materials and Methods

#### 5 *Generation of DC*

Immature DC were generated from peripheral blood monocyte precursors. Human PBMC from healthy donors, isolated through Ficoll-Hypaque density centrifugation were plated at  $1.5 \times 10^7$  per well in 6-well plates (Costar Corp., Cambridge, MA) in  
10 RPMI 1640 (Life Technologies, Paisley, Scotland) supplemented with 2mM glutamine, 100UI/ml penicillin and 10% FCS. After 2 h at  $37^{\circ}\text{C}$ , the non adherent cells were removed and the adherent cells were cultured in medium containing 500U/ml IL-4 (Pepro Tech Inc. Rocky Hill, NJ) and 800U/ml GM-CSF (kindly  
15 provided by Dr S. Osanto, LUMC, Leiden, NL) for a total of 7 days.

*Activation of immature DC with a CD8-CD40L fusion protein*  
Activation of DC through CD40 was performed with a fusion  
20 protein made of the extracellular domain of human CD40L and of the murine CD8a chain (CD8-CD40L). The CD8-CD40L cDNA described by Garrone et al (23) was transferred into an eukaryotic expression vector containing the hygromycin resistance gene, and used for the generation of stably  
25 transfected Chinese Hamster Ovary (CHO) cells. Culture supernatants containing the CD8-CD40L fusion protein were concentrated with a pressurized stirred cell system (Amicon, Inc., Beverly, MA), checked for binding to CD40 and tested for optimal DC activation conditions (not shown). DC were  
30 incubated at  $5 \times 10^5$ /ml/well in a 24-well plate (Costar Corp., Cambridge, MA) and activated in the presence of 1/10 CD8-CD40L supernatant. Cells and supernatants were analyzed after 48 h. Of note, control supernatants obtained from untransfected CHO cells or from CHO cells transfected with

the CD8a cDNA lacked DC activating functions and were similar to culture medium.

*DEX and RU486 treatment of DC*

- 5 Seven days immature DC were treated with  $10^{-6}$  M DEX (Sigma, St Louis, MO) in the presence of GM-CSF and IL-4 or GM-CSF alone. After 24 h, DC were analyzed or were further stimulated via CD40 by adding the CD8-CD40L fusion protein to the cultures as described above. In some experiments, the
- 10 glucocorticoid receptor antagonist RU485 (Roussel-UCLAF, Romainville, France) was used at 10mM final concentration, alone or in combination with DEX.

*Analysis of DC surface phenotype by flow cytometry*

- 15 Cells were stained on ice with FITC or PE-conjugated mouse monoclonal antibodies (MoAb) for 30 min in PBS 1% FCS and were analyzed on a FACScan<sup>®</sup> (Becton Dickinson, San Jose, CA). The following MoAb were used: FITC-anti-CD80 (BB1), PE-anti-CD86 (FUN-1), FITC-anti-CD40 (5C3), PE-anti-CD54 (HA 58) and
- 20 PE-anti-CD58 (1C3) (Pharmingen, San Diego, CA), PE-anti-CD14 (L243) and PE-anti-HLA-DR (Mf-P9) (Becton Dickinson), PE-anti-CD83 (HB15A) (Immunotech, Marseille, France) and PE-anti-HLA class I (Tu 149) (Caltag Laboratories, Burlingame, CA).

25

*Antigen uptake experiments*

- DC were resuspended in medium buffered with 25mM Hepes. FITC-BSA and FITC-mannosylated BSA (both from Sigma) were added at 1mg/ml final concentration and the cells were incubated at
- 30  $37^{\circ}\text{C}$ , or at  $0^{\circ}\text{C}$  to determine background uptake. After 1 h, DC were washed extensively with iced-cold PBS and analyzed by FACS<sup>®</sup> using propidium iodide to eliminate dead cells.

*Cytokine detection by ELISA*

Culture supernatants were analyzed in serial twofold dilutions in duplicate. IL-12p70 was detected using a solid phase sandwich ELISA kit (Diaclone Research, Besancon, France) (sensitivity 3pg/ml). For IL-12p40 and IFN- $\gamma$  detection, capture MoAb and polyclonal biotinylated detection Ab were obtained from Peter van de Meijde (BPRC, Rijswijk, NL) (sensitivity 10pg/ml). IL-10 was detected using the Pelikine compact human IL-10 ELISA kit (CLB, Amsterdam, NL) (sensitivity 3pg/ml).

*Allogeneic mixed lymphocyte reaction (MLR)*

Non adherent allogeneic adult PBMC from an unrelated individual were cultured in 96-well flat-bottom plates (Costar Corp., Cambridge, MA) at a density of  $1.5 \times 10^5$ /well with various numbers of g-irradiated (3,000 rads) DC, in triplicates. Proliferation was assessed on day 5 by [ $^3$ H]thymidine uptake (0.5mCi/well, specific activity 5Ci/mMol, Amersham Life Science, Buckinghamshire, UK) during a 16 h pulse.

*Th1 stimulation assays*

The Mycobacterium tuberculosis and M. leprae hsp65-specific, HLA-DR3-restricted CD4+ Th1 clone Rpl5 1-1 used in this study recognizes an hsp65 determinant corresponding to peptide residues 3 to 13 (p3-13) (24). HLA-DR-matched DEX-treated immature DC and their DEX-untreated counterparts were pulsed with 10mg/ml of p3-13 or with 10mg/ml of hsp65 for 2 h, washed extensively and stimulated through CD40 as described above. For Ag-pulsed DEX-treated immature DC, CD40 triggering was performed in the presence of DEX. Hsp65 specific T cells ( $10^4$ ) were cultured with different numbers of g-irradiated (3,000 rads) DC in 96-well flat-bottom plates (Costar Corp.) in triplicates for 3 days. [ $^3$ H]thymidine incorporation was measured on day 3 after a 16 h pulse. Before the addition of



[<sup>3</sup>H]thymidine, 50ml of supernatants were collected from each well and supernatants from triplicate wells were pooled to measure IFN-g production. To test hsp65-specific T cells responsiveness to a second potent antigenic challenge, 10<sup>4</sup> T cells were first cultured for 48h with 5 x 10<sup>3</sup> peptide-pulsed DC prepared as above, then harvested and allowed to rest in medium containing 5U/ml IL-2. Three days later, 10<sup>4</sup> viable T cells were restimulated with 5 x 10<sup>3</sup> peptide-pulsed DC generated from the same donor as used for the first culture and tested for their ability to proliferate and to produce IFN-g as previously described.

#### Statistical analysis

Covariance analysis was used to compare T cell proliferation and IFN-g production as a function of DC number, between DEX-treated CD40-triggered DC and DEX-untreated CD40-triggered DC (Fig. 5).

**Figure legends**

**Fig. 1** Pretreatment with DEX inhibits the phenotypic changes induced by CD40 ligation.

5 Seven days immature DC were cultured for 24h in the absence or the presence of  $10^{-6}$ M DEX and activated via CD40 with the CD8-CD40L fusion protein for 48h. The comparison with immature DC maintained in medium alone is shown. Empty histograms show the background staining with isotype controls

10 MoAb and solid histograms represent specific staining of the indicated cell surface markers. Specific mean fluorescence intensities are indicated. Mean fluorescence intensities of isotype controls were between 3 and 4. Data are representative of 4 independent experiments.

15

**Fig. 2** DC triggered through CD40 maintain an activated phenotype upon a subsequent DEX exposure. Immature DC were activated with the CD8-CD40L fusion protein. DEX ( $10^{-6}$ M) or medium control were added 48h later and cells

20 were analyzed after 2 additional days of culture. The comparison with immature DC maintained in medium alone is shown. Empty histograms show the background staining with isotype controls MoAb and solid histograms represent specific staining of the indicated cell surface markers. Specific mean

25 fluorescence intensities are indicated. Mean fluorescence intensities of isotype controls were between 3 and 5. Data are representative of 2 independent experiments.

**Fig. 3** Pretreatment with DEX does not affect the regulation of DC antigen uptake machinery.

30 Immature DC were incubated in the absence or the presence of  $10^{-6}$ M DEX for 24h and further activated or not via CD40 with the CD8-CD40L fusion protein for 48h. Cells were pulsed for 1h with medium containing either 1mg/ml FITC-BSA or 1mg/ml

35 FITC-mannosylated BSA. Empty histograms show the background

autofluorescence, Grey-filled histograms show the background uptake at 0°C and black-filled histograms show the specific uptake at 37°C. Data are representative of 3 independent experiments.

5

**Fig. 4** Pretreatment with DEX alters the cytokine secretion profile of CD40-triggered DC.

DEX-exposed or control immature DC were left in culture without further treatment or stimulated with the CD8-CD40L fusion protein. Culture supernatants were harvested 48h later and IL-10, IL-12p40 and IL-12p70 secretion were analyzed by specific ELISA. Data are representative from 6 independent experiments.

15 **Fig. 5** Pretreatment with DEX impairs the T cell stimulatory capacities of DC activated via CD40 and leads to a state of hyporesponsiveness of Th1 cells.

Allogeneic MLR: non adherent allogeneic PBMC were cultured with different numbers of CD40-triggered DC, DEX-treated CD40-triggered DC or immature DC. The proliferative response was measured on day 5.

20 Th1 stimulation assays: Hsp65-specific T cells were cultured with different numbers of HLA-DR matched CD40-triggered DC or with DEX-treated CD40-triggered DC pulsed with the hsp65

25 protein or with the specific p3-13 peptide epitope. The proliferative response and the T cell dependent IFN- $\gamma$  production were analyzed on day 3. Data are representative of 4 independent experiments.

30 **Fig. 6** DEX-treated DC triggered through CD40 induce a state of hyporesponsiveness in Th1 cells. Hsp65-specific T cells precultured with CD40-triggered DC or with DEX-treated CD40-triggered DC pulsed with the p3-13 peptide epitope were harvested after 48h, allowed to rest in the presence of 50U/ml  
35 IL2 for 3 days, and restimulated with p3-13-pulsed DC. The

proliferative response and IFN-g production were measured on day 3. Similar results were obtained in 2 independent experiments.

5 **Fig. 7** D1 cells were left untreated or cultured for two days in the presence of LPS (1 ug/ml) and/or DEX (10<sup>-6</sup> M). Surface expression of CD86 (B7.2), Class II MHC and CD40 was monitored by staining with the appropriate FITC-labeled Ab. and subsequent FACS-analysis. Similar results were obtained  
10 for D1 cells matured in the presence of murine CD40L (not shown).

**Fig. 8** Balb/c mice were immunised with 10<sup>-6</sup> D1 cells that were prepared as indicated. One week later, splenocytes were  
15 isolated and tested for their response against C57Bl/6 splenocytes in a mixed lymphocyte culture. After 24 h, culture supernatants were tested for IFN $\gamma$  content in a specific ELISA assay.

20 **Fig. 9** Balb/c mice (5 per group) were immunised as indicated. One week later, skin transplants of C57Bl/6 origin were grafted and graft survival was monitored.

REFERENCES

1. Banchereau J, Steinman RM: Dendritic cells and the control of immunity. *Nature* 392: 248, 1998
- 5 2. Sallusto F, Lanzavecchia A: Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony stimulating factor and downregulated by tumor necrosis factor  $\alpha$ . *J Exp Med* 179: 1109, 1994
- 10 3. Sallusto F, Cella M, Danieli C, Lanzavecchia A: Dendritic cells use macropinocytosis and the mannose receptor to concentrate macromolecules in the major histocompatibility complex class II compartment: downregulation by cytokines and bacterial products. *J Exp Med* 182: 389, 1995
- 15 4. Caux C, Massacrier C, Vandervliet B, Dubois B, van Kooten C, Durand I, Banchereau J: Activation of human dendritic cells through CD40 cross-linking. *J Exp Med* 180: 1263, 1994
5. Cella M, Sheidegger D, Palmer-Lehman K, Lane P, Lanzavecchia A, Alber G: Ligation of CD40 on dendritic cells triggers production of high levels of interleukin-12 and enhances T cell stimulatory capacity: T-T help via APC activation. *J Exp Med* 184: 747, 1996
- 20 6. Koch F, Stanzl U, Jennevin P, Janke K, Heufler C, Kampgen E, Romani N, Schuler G: High levels IL-12 production by murine dendritic cells: upregulation via MHC class II and CD40 molecules and downregulation by IL-4 and IL-10. *J Exp Med* 184: 741, 1996
- 25 7. Trinchieri G: Interleukin-12: a proinflammatory cytokine with immunoregulatory functions that bridge innate resistance and antigen-specific adaptive immunity. *Annu Rev Immunol* 13: 251, 1995
- 30 8. Kamanaka M, Yu P, Yasui T, Koshida K, Kawaba T, Horii T, Kishimoto T, Kikutani H: Protective role of CD40 in *Leishmania* major infection at two distinct phases of cell-mediated immunity. *Immunity* 4: 275, 1996
- 35

9. Yang Y, Wilson JM: CD40 ligand-dependent T cell activation: requirement of B7-CD28 signaling through CD40. *Science* 273: 1862, 1996
10. Mackey MF, Gunn JR, Ting P, Kikutani H, Dranoff G, Noelle RJ, Barth Jr RJ: Protective immunity induced by tumor vaccines requires interactions between CD40 and its ligand CD154. *Cancer Res* 57: 2569, 1997
11. Grewal IS, Foellmer HG, Grewal KD, Xu J, Hardardottir F, Baron JL, Janeway Jr CA, Flavell RA: Requirement for CD40 ligand in costimulation induction, T cell activation and experimental allergic encephalomyelitis. *Science* 273: 1864, 1996
12. Kirk AD, Harlan DM, Armstrong NN, Davis TA, Dong Y, Gray GS, Hong X, Thomas D, Fechner JH, Knechtle SJ: CTLA4-Ig and anti-CD40 ligand prevent renal allograft rejection in primates. *Proc Natl Acad Sci USA* 94: 8789, 1997
13. Durie FH, Aruffo A, Ledbetter J, Crassi KM, Green WR, Fast LD, Noelle RJ: Antibody to the ligand of CD40, gp39, blocks the occurrence of the acute and chronic forms of graft-vs-host disease. *J Clin Invest* 94: 1333, 1994
14. Almawi WY, Beyhum HN, Rahme AA, Rieder MJ: Regulation of cytokine and cytokine receptor expression by glucocorticoids. *J Leukoc Biol* 60: 563, 1996
15. Moser M, De Smedt T, Sornasse T, Tielemans F, and Chentoufi AA: Glucocorticoids down-regulate dendritic cell function in vitro and in vivo. *Eur J Immunol* 25: 2818, 1995
16. Kitajima T, Ariizumi K, Bergstresser PR, Takashima A: A novel mechanism of glucocorticoid-induced immune suppression: the inhibition of T cell-mediated terminal maturation of a murine dendritic cell line. *J Clin Invest* 98: 142, 1996
17. Viera PL, Kalinski P, Wierenga EA, Kapsenberg ML, de Jong EC: Glucocorticoids inhibit bioactive IL-12p70 production by in vitro-generated human dendritic cells without affecting their T cell stimulatory potential. *J Immunol* 161: 5245, 1998
18. Romani N, Gruner S, Brang D, Kampgen E, Lenz A, Trockenbacher B, Konwalinka G, Fritsch PO, Steinman RM,

- Schuler G: Proliferating dendritic cell progenitors in human blood. *J Exp Med* 180: 83, 1994
19. Randolph GC, Beaulieu S, Lebecque S, Steinman RM, Müller WA: Differentiation of monocytes into dendritic cells in a model of transendothelial trafficking. *Science* 282: 480, 1998
20. Rissoan MC, Soumelis V, Kadowaki N, Grouard G, Briere F, de Waal Malefyt R, Liu Y-J: Reciprocal control of T helper cell and dendritic cell differentiation. *Science* 283: 1183, 1999
21. Bender A, Sapp M, Schuler G, Steinman RM, Bhardwaj N: Improved methods for the generation of dendritic cells from non proliferating progenitors in human blood. *J Immunol Methods* 196:121, 1996
22. Romani N, Reider D, Heuer M, Ebner S, Kampgen E, Eibl B, Niederwieser D, Schuler G: Generation of mature dendritic cells from human blood: an improved method with special regard to clinical applicability. *J Immunol Methods* 196:137, 1996
23. Garrone P, Neidhardt E-M, Garcia E, Galibert L, van Kooten C, Banchereau J: Fas ligation induces apoptosis of CD40-activated human B lymphocytes. *J Exp Med* 182: 1265, 1995
24. Geluk A, van Meigaarden KE, Janson AAM, Drijfhout J-W, Meloen R, de Vries RRP, Ottenhoff T: Functional analysis of DR17(DR3)-restricted mycobacterial T cell epitopes reveals DR17 binding motif and enables the design of allele-specific competitor peptides. *J Immunol* 149: 2864, 1992
25. Engering AJ, Cella M, Fluitsma D, Brockhaus M, Hoefsmit ECM, Lanzavecchia A, Pieters J: The mannose receptor functions as a high capacity and broad specificity antigen receptor in human dendritic cells. *Eur J Immunol* 27: 2412, 1997
26. Tan MCAA, Mommaas AA, Drijfhout J-W, Jordens R, Onderwater JJM, Verwoerd D, Mulder AA, van der Heiden AN, Scheidegger D, Oomens LCJM, Ottenhoff THM, Tulp A, Neeffjes JJ, Koning F: Mannose receptor-mediated uptake of antigens strongly enhances HLA class II-restricted antigen

- presentation by cultured dendritic cells. *Eur J Immunol* 27: 2426, 1997
27. de Saint Vis B, Fugier-Vivier I, Massacrier C, Gaillhard C, Vanbervliet B, Ait-Yahia S, Banchereau J, Liu Y-J, Lebecque S, Caux C: The cytokine profile expressed by human dendritic cells is dependent on cell subtype and mode of activation. *J Immunol* 160: 1666, 1998
28. Piemonti L, Monti P, Allavena P, Sironi M, Soldini L, Leone BE, Socci C, Di Carlo V: Glucocorticoids affect human dendritic cell differentiation and maturation. *J Immunol* 162: 6473, 1999
29. Blotta MH, Dekruyff RH, Umetsu DT: Corticosteroids inhibit IL-12 production in human monocytes and enhance their capacity to induce IL-4 synthesis in CD4+ lymphocytes. *J Immunol* 158: 5589, 1997
30. Visser J, van Boxel-Dezaire A, Methrost D, Brunt T, de Kloet ER, Nagelkerken L: Differential regulation of interleukin 10 (IL-10) and IL-12 by glucocorticoids in vitro. *Blood*. 91: 4255, 1998
31. van Kooten C, Banchereau J: Function of CD40 on B cells, dendritic cells and other cells. *Curr Opin Immunol* 9: 330, 1997
32. Scheinman RI, P.C. Cogswell PC, Lofquist AK, and Baldwin Jr AS: Role of transcriptional activation of I kappa B alpha in mediation of immunosuppression by glucocorticoids. *Science* 270: 283, 1995
33. Auphan N, Dinato JA, Rosette C, Helmberg A, Karin N: Immunosuppression by glucocorticoids: inhibition of NF-kappa B activity through induction of I kappa B synthesis. *Science* 270: 286, 1995
34. Murphy TL, Cleveland MG, Kulesza P, Magram J, Murphy KM: Regulation of interleukin 12 p40 expression through an NF-kappa B half site. *Mol Cell Biol* 15: 5258, 1995
35. Yoshimoto T, Nagase H, Ishida T, Inoue J, Nariuchi H: Induction of interleukin-12 p40 transcript by CD40 ligation



- via activation of nuclear factor-kappaB. *Eur J Immunol* 27: 3461, 1997
36. Ridge JP, Di Rosa F, Matzinger P: A conditioned dendritic cell can be a temporal bridge between a CD4+ T-helper and a T-killer cell. *Nature* 393: 474, 1998
- 5 37. Bennett SR, Carbone FR, Karamalis F, Flavell RA, Miller JF, Heath WR: Help for cytotoxic-T-cell responses is mediated by CD40 signalling. *Nature* 393: 478, 1998
38. Schoenberger SP, Toes REM, van der Voort EI, Offringa R, Melief CJM: T-cell help for cytotoxic T lymphocytes is mediated by CD40-CD40L interactions. *Nature* 393: 480, 1998
- 10 39. Winzler C, Rovere P, Rescigno M, Granucci F, Penna G, Adorini L, Zimmermann VS, Davoust J, Ricciardi-Castagnoli P. Maturation stages of mouse dendritic cells in growth factor-dependent long-term cultures. *J Exp Med* 1997 Jan 20;185(2):317-28.
- 15 40. Schuurhuis DH, Laban S, Toes RE, Ricciardi-Castagnoli P, Kleijmeer MJ, van der Voort EI, Rea D, Offringa R, Geuze HJ, Melief CJ, Ossendorp F. Immature dendritic cells acquire CD8(+) cytotoxic T lymphocyte priming capacity upon activation by T helper cell-independent or -dependent stimuli. *J Exp Med* 2000 Jul 3;192(1):145-50
- 20

CLAIMS

1. A method for preparing a pharmaceutical composition for reducing an unwanted T cell response in a host, comprising culturing peripheral blood monocytes from said host to differentiate into dendritic cells, activating said dendritic cells in the presence of a glucocorticoid hormone and loading said dendritic cells with an antigen against which said T cell response is to be reduced.
2. A pharmaceutical composition for reducing an unwanted T cell response in a host, said composition being obtained by culturing peripheral blood monocytes from said host to differentiate into dendritic cells, activating said dendritic cells in the presence of a glucocorticoid hormone and loading said dendritic cells with an antigen against which said T cell response is to be reduced.
3. A method for reducing an unwanted T cell response in a host, comprising administering a composition of claim 2 to said host.
4. A method for reducing an unwanted T cell response in a host comprising culturing peripheral blood monocytes from said host to differentiate into dendritic cells, activating said dendritic cells in the presence of a glucocorticoid hormone and loading said dendritic cells with an antigen against which said T cell response is to be reduced and administering said composition to said host.
5. A method according to claim 1,3 or 4 whereby said activation is done through a CD40 receptor.
6. A method according to claim 5 whereby said activation involves incubation of the dendritic cells with either CD8-CD40L fusion protein, a trimeric form of CD40L consisting of CD40L-molecules to which a modified leucine zipper has been attached, anti-CD40 antibodies, or cells that express CD40L.
7. A method according to claim 5 whereby said activation involves incubation of the dendritic cells with lipopolysaccharide (LPS) or polyI/C.

8. A method according to claim 1, 3-7 whereby said dendritic cells are infected with one or more recombinant viruses encoding the antigen(s) of interest before activating said dendritic cells in the presence of a glucocorticoid hormone.
9. A method according to claim 1, 3-8 whereby said dendritic cells are incubated with one or more recombinant proteins or large (> 20 amino acids) synthetic peptides representing the antigen(s) of interest before activating said dendritic cells in the presence of a glucocorticoid hormone.
10. A method according to claim 1, 3 or 9 whereby said dendritic cells are incubated with cells or cell homogenate containing the antigen(s) of interest before activating said dendritic cells in the presence of a glucocorticoid hormone.
11. A method according to claim 1, 3-10 whereby said dendritic cells are loaded with synthetic peptides representing the antigen(s) of interest after activating said dendritic cells in the presence of a glucocorticoid hormone.
12. A method according to claim 1, 3-11 whereby said dendritic cells, after activation in the presence of a glucocorticoid hormone, secrete interleukin-10.
13. A method for obtaining a dendritic cell capable of tolerising a T-cell for an antigen comprising providing said dendritic cell with a glucocorticoid hormone, activating said dendritic cell and providing said dendritic cell with said antigen.
14. A method according to anyone of claims 1, 3-13, wherein said dendritic cell and/or a precursor thereof is provided with said glucocorticoid hormone in vitro.
15. A method according to anyone of claims 1, 3-14, wherein said T-cell is a T-helper cell.
16. An isolated dendritic cell prepared according to anyone of claims 1, 3-15 capable of functionally modifying an antigen-specific T-cell with respect to the response to said antigen.

17. A method for functionally modifying an antigen-specific T-cell comprising providing an dendritic cell according to claim 16 with said antigen and co-cultivating said T-cell and said dendritic cell.

5 18. A method according to claim 17, wherein said co-cultivating is performed in vitro.

19. A method according to claim 17 or claim 18, further comprising multiplying said functionally modified T-cell.

20. An isolated functionally modified T-cell obtainable by a  
10 method according to anyone of claims 17-19 that is capable, upon administration to the host, of reducing an unwanted immune response.

21. Use of a glucocorticoid hormone for obtaining an dendritic cell capable of functionally modifying a T-cell.

15 22. A pharmaceutical composition comprising an dendritic cell according to claim 16 and/or a functionally modified T-cell according to claim 20.

23. Use of an dendritic cell according to claim 16 and/or a functionally modified T-cell according to claim 20 for the  
20 preparation of a medicament.

24. A method for the treatment of an individual suffering from or at risk of suffering from a disease associated with at least part of the immune system of said individual comprising providing said individual with an dendritic cell  
25 according to claim 16 and/or a functionally modified T-cell according to claim 20.

25. A method according to claim 24, wherein said dendritic cell and or said T-cell is derived from an HLA-matched donor.

26. A method according to claim 24 or claim 25, wherein said  
30 dendritic cell and or said T-cell is derived from said individual.

27. Use of an dendritic cell according to claim 16 in a treatment for an individual suffering from an auto-immune disease, allergy, a graft versus host disease and/or a host  
35 versus graft disease.




Fig. 1




Fig. 2




Fig. 3




Fig. 4






Fig. 5




Fig. 6

Fig. 7. DEX modulates LPS-induced maturation of murine DC




Figure 8. Decreased allo-reactivity after pre-treatment with DEX-DC





Figure 9. Prolonged graft survival after pre-treatment with DEX-DC



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/NL 00/00712

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K39/00 A61K35/14 C12N5/08

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category \* Citation of document, with indication, where appropriate, of the relevant passages

Relevant to claim No.

X VIEIRA P L ET AL: "Glucocorticoids inhibit bioactive IL-12p70 production by in vitro-generated human dendritic cells without affecting their T cell stimulatory potential."  
JOURNAL OF IMMUNOLOGY,  
vol. 161, no. 10,  
15 November 1998 (1998-11-15), pages  
5245-5251, XP000973668  
Y cited in the application  
the whole document

1-7,  
9-16,  
22-27

8-12,  
14-16,  
22-27

---  
-/-

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claims or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*\* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*S\* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

7 February 2001

02.03.01

Name and mailing address of the ISA  
European Patent Office, P.O. Box 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer  
Bilang, J

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/NL 00/00712

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>LU L ET AL: "Adenoviral delivery of CTLA4lg into myeloid dendritic cells promotes their in vitro tolerogenicity and survival in allogeneic recipients" GENE THERAPY, MACMILLAN PRESS LTD., BASINGSTOKE, GB, vol. 6, no. 4, 1999, pages 554-563, XP000946114 ISSN: 0969-7128 the whole document</p>	<p>8-12, 14-16, 22-27</p>
A	<p>PIEMONTE L ET AL: "Glucocorticoids affect human dendritic cell differentiation and maturation." JOURNAL OF IMMUNOLOGY, vol. 162, no. 11, 1 June 1999 (1999-06-01), pages 6473-6481, XP002159692 cited in the application the whole document</p>	<p>1-27</p>
A	<p>GAO J X ET AL: "CD40-deficient dendritic cells producing interleukin-10, but not interleukin-12, induce T-cell hyporesponsiveness in vitro and prevent acute allograft rejection." IMMUNOLOGY, vol. 98, no. 2, October 1999 (1999-10), pages 159-170, XP000973593 the whole document</p>	<p>1-27</p>
A	<p>CHAKRABORTY ABALOKITA ET AL: "Stimulatory and inhibitory maturation of human macrophage-derived dendritic cells." PATHOBIOLOGY, vol. 67, no. 5-6, March 2000 (2000-03), pages 282-286, XP002159693 ISSN: 1015-2008 the whole document</p>	<p>1-27</p>
P,X	<p>REA D ET AL: "Glucocorticoids transform CD40-triggering of dendritic cells into an alternative activation pathway resulting in antigen-presenting cells that secrete IL-10." BLOOD, vol. 95, no. 10, 15 May 2000 (2000-05-15), pages 3162-3167, XP002159694 the whole document</p>	<p>1-27</p>

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/NL 00/00712**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claim(s) 3-12, 14, 16, 17, 19, 24, and 27 are directed to a diagnostic method practised on the human/animal body or treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.